



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**331. Proposed by T. H. GRONWALL, Chicago, Illinois.**

To show that:

- (1)  $\frac{d^n}{dx^n} \left( \frac{\sin x}{x} \right) = \frac{1}{x^{n+1}} \int_0^x y^n \sin \left( y + \frac{n+1}{2} \pi \right) dy,$   
 (2)  $\frac{d^n}{dx^n} \left( \frac{1 - \cos x}{x} \right) = \frac{1}{x^{n+1}} \int_0^x y^n \sin \left( y + \frac{n}{2} \pi \right) dy.$

SOLUTION BY ELIJAH SWIFT, Princeton, N. J.

Assume that the formula is true for  $n - 1$ .

Then

$$\frac{d^{n-1}}{dx^{n-1}} \left( \frac{\sin x}{x} \right) = \frac{1}{x^n} \int_0^x y^{n-1} \sin \left( y + \frac{n}{2} \pi \right) dy.$$

Differentiating both sides and using the formula we have to prove

$$(A) \quad \frac{1}{x^{n+1}} \int_0^x y^n \sin \left( y + \frac{n+1}{2} \pi \right) dy = -\frac{n}{x^{n+1}} \int_0^x y^{n-1} \sin \left( y + \frac{n}{2} \pi \right) dy \\ + \frac{1}{x^n} \cdot x^{n-1} \sin \left( x + \frac{n}{2} \pi \right).$$

or

$$(B) \quad \int_0^x y^n \sin \left( y + \frac{n+1}{2} \pi \right) dy = -n \int_0^x y^{n-1} \sin \left( y + \frac{n\pi}{2} \right) dy \\ + x^n \sin \left( x + \frac{n}{2} \pi \right).$$

Differentiating  $B$  as to  $x$ ,

$$(C) \quad x^n \sin \left( x + \frac{n+1}{2} \pi \right) = -nx^{n-1} \sin \left( x + \frac{n\pi}{2} \right) + nx^{n-1} \sin \left( x + \frac{n\pi}{2} \right) \\ + x^n \cos \left( x + \frac{n}{2} \pi \right),$$

which is obviously an identity.

Hence  $(B)$  is an identity, except for an additive constant. This constant must be zero, as we see if we let  $x$  approach zero.  $(A)$  follows from  $(B)$ . Then since the given formula holds for  $n = 0$ , it holds for all values of  $n$ .

Formula (2) may be derived in the same manner.

Also solved by A. M. Harding, W. C. Eells and the Proposer.

## NEWS AND NOTES.

FLORIAN CAJORI, CHAIRMAN OF THE COMMITTEE.

The March number of *Popular Science Monthly* contains an article on "Henri Poincaré as an Investigator" by PROFESSOR JAMES B. SHAW.

PROFESSOR G. A. MILLER, of the University of Illinois, will teach in the summer session of the University of California.